

Scattering and Thermodynamics of Fractionally-Charged Supersymmetric Solitons

P. Fendley

Department of Physics
Boston University
590 Commonwealth Avenue
Boston, MA 02215

K. Intriligator

Lyman Laboratory of Physics
Harvard University
Cambridge, MA 02138

We show that there are solitons with fractional fermion number in integrable $N=2$ supersymmetric models. We obtain the soliton S-matrix for the minimal, $N=2$ supersymmetric theory perturbed in the least relevant chiral primary field, the $\Phi_{(1,3)}$ superfield. The perturbed theory has a nice Landau-Ginzburg description with a Chebyshev polynomial superpotential. We show that the S-matrix is a tensor product of an associated ordinary ADE minimal model S-matrix with a supersymmetric part. We calculate the ground-state energy in these theories and in the analogous $N=1$ case and $SU(2)$ coset models. In all cases, the ultraviolet limit is in agreement with the conformal field theory.

1. Introduction

Integrable models have the striking property that in a collision all momenta are conserved individually, and that the n -body S-matrix factorizes into a product of two-body ones. The enormous number of constraints this implies means that the exact S-matrix can often be conjectured [1,2]. The exact S-matrix encodes, nonperturbatively, physical information about our theory; it is just a matter of extracting the information. One way to extract physical information from the exact S-matrix of a 1+1 dimensional integrable theory is to do thermodynamics via the thermodynamic Bethe ansatz [3,4]. If the S-matrix is exact, the results of the TBA are non-perturbative. For massive, integrable theories obtained from perturbing a conformal theory by a relevant operator, for instance, we can compare the Casimir energy predicted by the TBA with the Casimir energy as calculated in the perturbed conformal theory. Thus an important consistency check on a conjectured scattering theory is to see if the Casimir energy goes over to the correct central charge in the UV limit (where the perturbation goes away). The next corrections to the Casimir energy can also be computed from the conformal theory and compared with that predicted from the scattering theory. In this way, the S-matrix contains a tremendous amount of information about the theory. Some excited-state energies can also be obtained in this manner [5-7].

Consider, for example, perturbing the p -th unitary minimal ($N=0$) conformal theory by the least relevant operator, the $\Phi_{(1,3)}$ field:

$$S_p \rightarrow S_p + \lambda \int \Phi_{(1,3)} d^2x. \quad (1.1)$$

Perturbative evidence suggests that this theory has an infinite number of conserved currents [8]. It was explicitly shown that there is at least one kinematic conserved current (beyond the energy-momentum tensor), which insures that there must be completely elastic and factorizable scattering. For $\lambda > 0$ the perturbed theory (1.1) gives a RG trajectory flowing from the p -th minimal model in the UV limit to the $(p-1)$ -th minimal model in the IR limit. Since there is a non-trivial IR limit, there must be massless particles in the spectrum; this situation is discussed in [9]. For $\lambda < 0$ the perturbed theory (1.1) is massive and the integrability requires factorized scattering. The exact scattering theory associated with this perturbed theory was conjectured in [10-12]. The ground-state energy associated with this conjectured scattering theory was considered using the thermodynamic Bethe ansatz

in [13] and [12]. It was verified to the first few orders in the UV limit ($\lambda \rightarrow 0$) that these results agree with those obtained from perturbing the conformal theory.

In this paper we will propose a scattering theory for the $N=2$ supersymmetric analog of (1.1), i.e. the $N=2$ supersymmetric minimal models perturbed in the least relevant operator, the $\Phi_{(1,3)}$ superfield. We assume integrability for these theories—this has been shown to lowest order in perturbation theory, and it is argued that the supersymmetry prevents any corrections [14]. These perturbed theories have a nice Landau-Ginzburg description [15] with a Chebyshev polynomial as the perturbed superpotential.

The Landau-Ginzburg analysis indicates that these $N=2$ theories are closely related to their $N=0$ analog scattering theories (1.1). In fact, we will show that the S-matrix for the $N=2$ theory is a direct product of the S-matrix for the $N=0$ analog (1.1) with a basic $N=2$ supersymmetric S-matrix. This structure was conjectured in [16], as the simplest S-matrix consistent with the quantum-group symmetry of the model. Taking the nonrenormalization of the superpotential to hold even nonperturbatively, an assertion which has been supported in many ways, we know the exact superpotential and the exact soliton spectrum, including the masses of the solitons [17]. Thus we show that the solitons conjectured in [16] do exist. However, we have a very different interpretation of the supersymmetric part of the S-matrix—in [16] the supersymmetry is postulated to act nonlocally, whereas here it acts locally.

The striking feature of a Chebyshev superpotential is that it has a set of degenerate wells with equal-mass solitons interpolating between adjacent ones. The bosonic part is the type of $N=0$ Landau-Ginzburg potential expected for theory (1.1)[10]¹. This provides a nice way of thinking about the Landau-Ginzburg potential for the $N=0$ minimal model. It also indicates why the soliton spectrum of the $N=2$ theory has its tensor product structure: the bosonic $N=0$ solitons become a supermultiplet in the supersymmetric theory. (This tensor-product structure has also been conjectured for the $N=1$ case [19,20].) We will show that this behavior generalizes to the $N=2$ D -series and to the exceptional invariants. In fact, all the $N=2$ theories we discuss have a close relation to an $N=0$ model. The Landau-Ginzburg soliton picture has been discussed for many $N=2$ theories [21], so making this correspondence can provide intuition concerning many non-supersymmetric models.

One fascinating feature we find is that the solitons have fractional fermion number. This phenomena was first observed some time ago [22] in a two-dimensional ϕ^4 theory

¹ This degenerate-well potential can also be predicted by studying the associated lattice model [18].

coupled to a Dirac fermion. In the Landau-Ginzburg picture, our models have the same structure as that treated in [22]. However, our models are exactly soluble and we obtain the exact S-matrices. The models discussed in this paper have solitons with charges of $\pm\frac{1}{2}$. In our next paper [23], we will find the S-matrix for the $N=2$ minimal models perturbed by the most-relevant operator, as well as for models with generalized Chebyshev potentials. These theories have more general (but still rational) fractional charges. In particular, the charges are multiples of $1/(k+1)$ in the most-relevant perturbation of the k -th minimal model.

We will provide checks on the $N=2$ scattering theories by verifying that the Casimir energy obtained from the TBA goes over to the correct central charges in the UV limit. We will also perform this analysis for the analog $N=1$ discrete series [19,20] and for the perturbed $SU(2)$ coset models[20], since these fit nicely into our general picture. We will also calculate $\text{tr}(-1)^F$ in the $N=2$ theories as a further check on the TBA equations.

2. N=2 soliton structure

Consider a $N=2$ theory with an effective LG description characterized by some superpotential $W(X)$ (X is a chiral superfield consisting of a complex boson and a Dirac fermion). Since the bosonic part of the potential is $|W'(X)|^2$, the vacua are the points in the complex X plane where $dW = 0$. The solitons X_{ij} are the finite energy solutions to the equations of motion connecting the i -th and j -th vacua: $X(\sigma = -\infty) = X^{(i)}$, $X(\sigma = +\infty) = X^{(j)}$ (as discussed in [17], not all such kinks are to be regarded as fundamental solitons). In the soliton sector corresponding to a soliton X_{ij} with mass m and rapidity θ , the $N=2$ superalgebra is [24,17]:

$$\begin{aligned} Q_+^2 &= Q_-^2 = \overline{Q}_+^2 = \overline{Q}_-^2 = \{Q_+, \overline{Q}_-\} = \{Q_-, \overline{Q}_+\} = 0 \\ \{Q_+, \overline{Q}_+\} &= 2\Delta W & \{Q_-, \overline{Q}_-\} &= 2(\Delta W)^* \\ \{Q_+, Q_-\} &= 2me^\theta & \{\overline{Q}_+, \overline{Q}_-\} &= 2me^{-\theta}, \end{aligned} \quad (2.1)$$

where $\Delta W = W(X^{(j)}) - W(X^{(i)})$. It follows from the above algebra that there is a Bogomolny mass bound $m \geq |\Delta W|$ [24,17]. The basic supermultiplet irreducible representation of (2.1) for solitons saturating this mass bound (the only type of solitons which we consider) is a doublet consisting of solitons $u(\theta)$ and $d(\theta)$ with:

$$\begin{aligned} Q^-|u(\theta)\rangle &= \sqrt{2m}e^{\theta/2}|d(\theta)\rangle & \overline{Q}^+|u(\theta)\rangle &= \omega\sqrt{2m}e^{-\theta/2}|d(\theta)\rangle \\ Q^+|d(\theta)\rangle &= \sqrt{2m}e^{\theta/2}|u(\theta)\rangle & \overline{Q}^-|d(\theta)\rangle &= \omega^*\sqrt{2m}e^{-\theta/2}|u(\theta)\rangle, \end{aligned} \quad (2.2)$$

where $\omega = \Delta W/m$. All other actions annihilate the states.

Our theory has a conserved $U(1)$ charge F corresponding to fermion number. The generators $Q^\pm$ have fermion number ± 1 , whereas $\overline{Q}^\pm$ have fermion number ∓ 1 (this notation is less bizarre at the conformal point where left and right fermion number are separately conserved). In soliton sectors the fermion number operator F generally picks up an additive constant piece, leading to fractional fermion number [22,25,26]. In fact, it can be shown via adiabatic or index theorem techniques [25,26] that the fractional part of the fermion number in a soliton sector X_{ij} of our theory is given by

$$f = -\frac{1}{2\pi} (Im \ln W''(X)) \Big|_{X^{(i)}}^{X^{(j)}}. \quad (2.3)$$

In our soliton doublet, $u(\theta)$ has fermion number e and $d(\theta)$ has fermion number $e - 1$, explaining why they haven't been labeled as boson and fermion. We will often label the solitons by their fermion number. The phenomena of fractional fermion number in 1+1 dimensions occurs in physical polymer systems where one extra bond (fermion) can be distorted into n solitons and, thus, each soliton carries fermion number $1/n$ [27].

The supersymmetry is defined on multi-particle states in the usual manner. Since Q is fermionic, one picks up phases when Q is brought through a particle with fermion number. For example, bringing Q through a fermion results in a minus sign. Since we will have fractional charges, we must generalize this notion to

$$Q^\pm |e_1 e_2\rangle = |(Q^\pm e_1) e_2\rangle + e^{\pm i\pi e_1} |e_1 (Q^\pm e_2)\rangle, \quad (2.4)$$

where the action on one soliton ($Q^\pm e$) is defined as in (2.2). The charges $\overline{Q}^\mp$ act with the same phases as $Q^\pm$. In notation analogous to that used for coproducts in a quantum group, (2.4) reads

$$\Delta(Q^\pm) = Q^\pm \otimes 1 + e^{\pm i\pi F} \otimes Q^\pm, \quad (2.5)$$

where F is the fermion-number operator. This similarity with the quantum-group action is not a coincidence, since $N=2$ supersymmetry is a special case of a quantum group [16].

The fractional fermion number is crucial for obtaining the correct soliton content and S-matrices. In [17], for example, the fractional charge structure was ignored. It was thus mistakenly conjectured (in the context of minimal models perturbed in the most relevant operator) on the basis of CPT that each soliton supermultiplet should be a quadruplet—a tensor product of two copies of the basic soliton doublet discussed above. Taking proper account of the fractional fermion number, one sees that this doubling is unnecessary.

Corresponding to this doubling, the S-matrix obtained in [17] is the tensor product of the correct S-matrix with its complex conjugate. The S-matrices for this class of integrable $N=2$ theories and the thermodynamic calculations (which confirm the S-matrices) will be discussed in [23]. In this paper we focus on another (simpler) class of integrable $N=2$ theories, $N=2$ minimal models perturbed in the *least* relevant chiral primary field.

3. Chebyshev Superpotentials

There is considerable evidence that the $N=2$ minimal models perturbed in the least relevant chiral primary field (the $\Phi_{(1,3)}$ perturbation) are integrable [14]. Using the results of [28], the effective superpotentials characterizing these perturbed theories are given by Chebyshev polynomials ²:

$$W_{k+2}(X = 2 \cos \theta) = \frac{2 \cos(k+2)\theta}{k+2}, \quad (3.1)$$

so that $W(X) = X^{k+2}/(k+2) - X^k + \dots$. For convenience, we have set the perturbing parameter to one (the powers of this parameter can be put back in by charge counting; e.g., $W_5 = X^5/5 - \beta X^3 + \beta^2 X$). These perturbed theories are intimately connected with $SU(2)_k$. The chiral ring structure constants (in the natural basis [28]), for example, are the fusion rules of $SU(2)_k$ [30].

The vacua of our theory (3.1) are the $k+1$ solutions of $dW(X) = \sin(k+2)\theta / \sin \theta = 0$:

$$X^{(r)} = 2 \cos \frac{\pi r}{k+2} \quad r = 1, \dots, k+1. \quad (3.2)$$

Thus all the vacua are on a line in the X -plane. This $(k+1)$ -well potential is characteristic of the $N=0$ analog theories (1.1) as well—the main difference here is that we have also the fermions. Using the methods of [17], it is easy to show that there exists a fundamental soliton connecting each of the adjacent critical points (3.2). Thus the spectrum consists of k solitons $X_{r(r+1)}$ for $r = 1, \dots, k$ and their k antisolitons $X_{(r+1)r}$. Any other possible soliton will break apart into two or more of these solitons. Taking our solitons to saturate the mass bound $m \geq |\Delta W|$, using the value of the superpotential (3.1) at the critical points

$$W(X^{(r)}) = \frac{2(-1)^r}{k+2}, \quad (3.3)$$

² The fact that our integrable perturbation should be in one of the flat directions of [28] is supported by [29]. Our results can be viewed as additional confirmation.

we see that our fundamental solitons connecting adjacent vacua all have equal mass $m = |\Delta W| = 4/(k+2)$.

Each of these solitons is a supermultiplet: the doublet discussed in the last section. From (2.3), we find that $u_{r(r+1)}(\theta)$ has fermion number $1/2$ and soliton $d_{r(r+1)}(\theta)$ has fermion number $-1/2$. The corresponding $2k$ antisolitons have opposite fermion numbers. For the $k=1$ case this structure is just that discussed in [22]: the Dirac equation has one zero mode in the presence of a soliton. (The supersymmetry requires that there be at least one; it turns out that there is only one.) When quantized, this Dirac zero mode results in a doublet of states, and the charge-conjugation symmetry requires that they have charge $\pm 1/2$.

Our soliton spectrum of $4k$ particles, consisting of a two-dimensional supermultiplet (2.2) for each soliton connecting adjacent vacua, is exactly the structure conjectured in [16] for these models; although, as we will detail in sect. 4, the action of supersymmetry is different. We will find an S-matrix for these solitons in the next section, and in sect. 6 we will show that the TBA calculation gives the expected central charges in the conformal limit. If one ignores this supermultiplet structure, one obtains exactly the soliton structure conjectured [10,11] (and verified to a large extent [13]) for the $N=0$ minimal models (1.1).

3.1. *D-series, E_6 , E_7*

The theories (3.1) correspond to perturbing the A_{k+1} minimal models in the least relevant operator. We found that the vacua of the theory correspond to the nodes of the A_{k+1} Dynkin diagram and the fundamental solitons correspond to the lines in the Dynkin diagram connecting the nodes. There are more $N=2$ minimal models, corresponding to the *D-series* and the exceptional models E_6 , E_7 and E_8 [15]. The *D-series* should be integrable when perturbed by the least-relevant operator, because each model is an orbifold of an *A-series* model treated in the previous subsection, and the conserved charge discussed in [14] is invariant under orbifolding. One expects that the exceptional models perturbed by their least-relevant operator to be integrable as well. The effective Landau-Ginzburg potentials for these perturbed theories are again obtained using the results of [28]. For the *D-series*, E_6 and E_7 these potentials were obtained explicitly in [31]:

$$\begin{aligned} W_{D_k}(x, y) &= \frac{1}{k-1} \cos(k-1)\phi + \frac{(-1)^{k-1}}{2} xy^2 \quad \text{where } 2 \cos \phi = 2 - x \\ W_{E_6}(x, y) &= \frac{x^3}{3} + \frac{y^4}{4} - xy^2 + \frac{y^2}{2} - \frac{x}{12} \\ W_{E_7}(x, y) &= \frac{x^3}{3} + \frac{xy^3}{3} - x^2y + \frac{4x^2}{9} - \frac{xy}{9} + \frac{x}{81} + \frac{1}{4374}, \end{aligned} \tag{3.4}$$

where, again, we have scaled the perturbing parameter to one. Minimizing these potentials, one finds that the vacua of these theories correspond to the nodes of the relevant Dynkin diagrams. We conjecture that the fundamental solitons are the lines in the Dynkin diagrams connecting the nodes. Just as in the A_{k+1} case, W alternates in sign between connected nodes and, thus, all solitons have equal mass. The $N=2$ structure again requires each soliton to consist of a (u, d) doublet with charges $\pm 1/2$.

4. The S-matrices

4.1. $X^3/3 - \beta X$ (sine-Gordon at its $N=2$ point)

The perturbed theory (3.1) with $k = 1$ corresponds to the sine-Gordon model, with coupling at the $N=2$ point: $\beta^2 = \frac{2}{3}8\pi$ in the conventional normalization. The $N=2$ symmetry is a special case of the general quantum group symmetry of the sine-Gordon model at any coupling [32,33]. Even though the $X^3/3 - \beta X$ model is the “same” as the sine-Gordon at this coupling, we find the action of supersymmetry in the two descriptions is very different. The difference arises from the fact that sine-Gordon and the $X^3/3 - \beta X$ are different local projections of the same theory [34]. This is analogous to the Ising model, where the spectrum can be either a free fermion or a strongly-interacting boson with S-matrix $S = -1$. They can be mapped on to each other by the non-local Jordan-Wigner transformation, but they most definitely are not identical. The same type of behavior happens in our $N=2$ model. The standard sine-Gordon S-matrix at this point [1] does not have an obvious supersymmetry: it must be realized nonlocally on the sine-Gordon solitons [10,16]. Even though the S-matrix in this section is formally the same as that of sine-Gordon at the appropriate coupling, ours does in fact allow a local action of supersymmetry. It is likely that there is an analog of the Jordan-Wigner transformation for this model, but we have not found one.

Fermion-number conservation means that the S-matrix for the process $|J(\theta_1)\overline{K}(\theta_2)\rangle \rightarrow |L(\theta_2)\overline{M}(\theta_1)\rangle$ (corresponding to a soliton X_{12} colliding with a soliton X_{21}) is of the form

$$\begin{array}{cc} u\overline{u} & d\overline{d} \\ u\overline{u} & d\overline{d} \end{array} \begin{pmatrix} c & b \\ b & c \end{pmatrix} \qquad \begin{array}{cc} u\overline{d} & d\overline{u} \\ d\overline{u} & u\overline{d} \end{array} \begin{pmatrix} 0 & a \\ a & 0 \end{pmatrix}, \quad (4.1)$$

where, again, u and d have charges $\pm 1/2$ respectively (and their antiparticles have opposite charges). Demanding that the S-matrix commutes with the supersymmetry generators with the action (2.2) and (2.5) requires the S-matrix elements to be

$$\begin{aligned} a &= Z(\theta) \cosh \frac{\theta}{2} \\ b &= Z(\theta) i \sinh \frac{\theta}{2} \\ c &= Z(\theta). \end{aligned} \tag{4.2}$$

Crossing, unitarity and the stipulation that there be no extra bound states fixes $Z(\theta)$ to be

$$\begin{aligned} Z(\theta) &= \frac{1}{\cosh \frac{\theta}{2}} \prod_{j=1}^{\infty} \frac{\Gamma^2(-\frac{\theta}{2\pi i} + j) \Gamma(\frac{\theta}{2\pi i} + j + \frac{1}{2}) \Gamma(\frac{\theta}{2\pi i} + j - \frac{1}{2})}{\Gamma^2(\frac{\theta}{2\pi i} + j) \Gamma(-\frac{\theta}{2\pi i} + j + \frac{1}{2}) \Gamma(-\frac{\theta}{2\pi i} + j - \frac{1}{2})} \\ &= \frac{1}{\cosh \frac{\theta}{2}} \exp\left(\frac{i}{4} \int_{-\infty}^{\infty} \frac{dt}{t} \frac{\sin \theta}{\cosh^2 \frac{\pi t}{2}}\right). \end{aligned} \tag{4.3}$$

It is crucial in this derivation that $\Delta W/m$ is 1 for the X_{12} and -1 for the X_{21} . Tensoring the above S-matrix with itself gives the S-matrix obtained in [17] where, as discussed in the last section, the basic soliton supermultiplet was unnecessarily doubled. Our thermodynamic calculations will confirm the above S-matrix.

This S-matrix is the same as that of the sine-Gordon model at $\beta^2 = \frac{2}{3}8\pi$ if we identify u and $\bar{d}$ with the soliton of that model and d and $\bar{u}$ with the antisoliton; the fermion number in the local description becomes the topological charge in sine-Gordon. For purposes of the ground-state thermodynamics done in sect. 6, this distinction is irrelevant; both S-matrices have the same TBA system and hence the same Casimir energy. However, we emphasize that the interpretations of the action of supersymmetry are very different.

4.2. $N=2$ minimal models with $\Phi_{1,3}$ perturbation

As discussed in sect. 3.1, the soliton structure resulting from the Chebyshev superpotential is that of the corresponding $N=0$ minimal model with additional $N=2$ structure. It does not immediately follow that the S-matrices are a direct product of the $N=0$ soliton S-matrix and an $N=2$ part, because the representation of the supersymmetry algebra depends on $\Delta W/m$, and this can depend on which solitons are being scattered. For this reason, the S-matrices for the $N=2$ minimal models perturbed by the most relevant operator (to be discussed in [23]) cannot be a tensor product. However, with the $\Phi_{1,3}$ (least-relevant) perturbation, $\Delta W/m$ just alternates between ± 1 (i.e. $\Delta W/m = (-1)^r$ for

$X_{r(r+1)})$ and, thus, all of the two-dimensional supermultiplets obey the same supersymmetry algebra as the $X^3 - \beta X$ model discussed in the previous section. From the point of view of the supersymmetry, every scattering process for all Chebyshev superpotentials looks the same. Thus the S-matrix for (3.1) is a direct product:

$$S_k^{N=2}(\theta) = S_k^{N=0}(\theta) \otimes S_{k=1}^{N=2}(\theta), \quad (4.4)$$

where $S_k^{N=0}$ is the S-matrix found in [10,11] for the massive scattering theory coming from the $N=0$ minimal model $SU(2)_1 \otimes SU(2)_k / SU(2)_{k+1}$ perturbed as in (1.1), and the S-matrix $S_{k=1}^{N=2}$ for the supersymmetric part is the one discussed in the previous subsection. This direct-product structure (4.4) was found in [19,20] for the $N=1$ supersymmetric case and in [16] for the $N=2$ case. Again, the S-matrix (4.1) allows a local action of supersymmetry, whereas the sine-Gordon S-matrix allows a nonlocal action. Both choices give the same thermodynamics. The nice thing about the LG analysis is that it gives the Chebyshev potential and hence the soliton structure directly.

The D_k series, E_6 and E_7 have the same tensor-product structure, and the $N=2$ part is exactly the same:

$$S_G^{N=2}(\theta) = S_G^{N=0}(\theta) \otimes S_{k=1}^{N=2}(\theta) \quad (4.5)$$

where $S_{k=1}^{N=2}$ is the basic $N=2$ S-matrix, obtained in this section, describing how the supermultiplet elements u and d scatter. The $N=0$ part $S_G^{N=0}$ is easy to find by using the correspondence between a factorizable S-matrix and the Boltzmann weights for a lattice model. In models with solitons, each vacuum corresponds to a “height” in an RSOS lattice model; for example, the $N=0$ A_n model S-matrix is equal to the Boltzmann weights for the critical RSOS lattice model A_{n-1} [18]. We conjecture that the D_n and E S-matrices are the Boltzmann weights for the critical RSOS models associated with the D_n and E Dynkin diagrams [36].

5. Thermodynamic Bethe Ansatz

The TBA allows us to extract the Casimir energy for our theory on a circle of length R . The only required input are the soliton masses and their two-body elastic S-matrix. The basic idea is to compute the minimum free energy of a system of solitons at temperature $T \equiv 1/R$ living on a circle of length $L \rightarrow \infty$. This is accomplished by finding the allowed energy levels on the circle and then filling the levels so as to minimize the free energy

F . The TBA analysis is thus in the grand canonical picture, with a chemical potential μ [4,37] (general chemical potentials enter in the calculation of excited-state energies[7]). Our system can be viewed as a two-dimensional system on a torus with length L corresponding to the compactified space and length $R = 1/T$ corresponding to compactified time. It thus follows that the ground-state energy $E(R)$ of our system on a circle of periodicity R is given in terms of our minimum free energy by

$$E(R) = RF_{min}(R, L)/L \quad \text{as } L \rightarrow \infty, \quad (5.1)$$

when $\mu = 0$.

Consider N particles with rapidities $\theta_1, \dots, \theta_N$ on the circle. The allowed wavefunction must be invariant upon bringing every rapidity θ_k around the circle, through the others and back to its starting point. In other words we must have

$$e^{im_k \sinh \theta_k L} T(\theta_k | \theta_{k+1}, \dots, \theta_N, \theta_1, \dots, \theta_{k-1}) \psi = \psi, \quad (5.2)$$

where $T(\theta_k | \theta_{k+1}, \dots, \theta_{k-1})$ is the transfer matrix for bringing particles with rapidity θ_k through the others. Our allowed wavefunctions must satisfy (5.2) for every allowed rapidity. In other words, the allowed wavefunctions must be simultaneous eigenvectors of the transfer matrices for bringing each rapidity through the others and the eigenvalues must satisfy (5.2) (the Yang-Baxter relation ensures that the transfer matrices commute). This condition quantizes the allowed particle energies $E_k = m_k \cosh \theta_k$ on the circle. We thus need to know how to diagonalize the transfer matrix in order to even write down the TBA system of equations. It is for this reason that the TBA equations have been derived for only a few theories with non-diagonal S-matrices [13,35]. In the next subsection we will review the warmup case where the S-matrix is diagonal. In the following subsection we will discuss some aspects of our case where the S-matrix is not diagonal.

5.1. The TBA equations for diagonal S-matrices

When the S-matrix is diagonal, i.e. the only scattering is of the type $a(\theta_1)b(\theta_2) \rightarrow S_{ab}(\theta_1 - \theta_2)b(\theta_2)a(\theta_1)$, the transfer matrix in (5.2) is diagonal for the N particle wavefunction describing particles of rapidity θ_i with the i -th particle having definite species a_i . The single-valuedness condition (5.2) then becomes simply:

$$e^{im_k \sinh \theta_k L} \prod_{j \neq k} S_{a_k a_j}(\theta_k - \theta_j) = 1, \quad (5.3)$$

an interacting-model generalization of the one particle relation $p_k = 2\pi n_k/L$. In the large L limit (5.3) gives the distribution $P_a(\theta)$ of allowed rapidity levels for particles of species a in terms of the distributions $\rho_b(\theta')$ of rapidity levels actually occupied by particles of species b :

$$2\pi P_a(\theta) = mL \cosh(\theta) + \sum_b \int d\theta' \rho_b(\theta') \phi_{ab}(\theta - \theta'), \quad (5.4)$$

where

$$\phi_{ab}(\theta) \equiv -i \frac{\partial \ln S_{ab}(\theta)}{\partial \theta}.$$

It is convenient to define quantities $\epsilon_a(\theta)$ by

$$\frac{\rho_a(\theta)}{P_a(\theta)} = \frac{\lambda_a e^{-\epsilon_a(\theta)}}{1 + \lambda_a e^{-\epsilon_a(\theta)}}, \quad (5.5)$$

where the fugacity $\lambda_a = e^{\mu_a R}$. This definition reflects the fact that all particle species in this paper (bosons and fermions) have $S_{aa}(0) = -1$ and, thus, at most one particle of a given species occupies a given momentum level³.

The distributions $\rho_a(\theta)$ are now chosen so as to minimize the free energy. This yields the TBA integral equations for the $\epsilon_a(\theta)$ in the presence of a general chemical potential [4,37]:

$$\epsilon_a(\theta) = m_a R \cosh(\theta) - \sum_b \int \frac{d\theta'}{2\pi} \phi_{ab}(\theta - \theta') \ln(1 + \lambda_b e^{-\epsilon_b(\theta')}). \quad (5.6)$$

Using (5.1) it can then be shown that the ground-state energy of the system can be written in terms of the ϵ_a as:

$$E_\lambda(R) = - \sum_a \frac{m_a}{2\pi} \int d\theta \cosh \theta \ln(1 + \lambda_a e^{-\epsilon_a(\theta)}). \quad (5.7)$$

Having obtained the system (5.6) appropriate for a given scattering theory, one can compute the Casimir energy $E(R)$ by (numerically) solving this set of coupled integral equations. The UV limit ($m \rightarrow 0$) and the IR limit ($m \rightarrow \infty$) of the Casimir energy can be obtained analytically and are discussed in sects. 5.3 and 5.4 respectively.


³ We define the S-matrix elements so that free fermions scattering off of one another have $S=-1$: this corresponds to ordering the creation operators by a particle's position in space. This definition enables us to use the Yang-Baxter equation and do the thermodynamics ignoring the fact that there are particles with non-zero fermion number.

5.2. Obtaining TBA Equations for non-diagonal S-matrices

We will describe the transfer matrices in (5.2) (one matrix for each particle k) in terms of a single transfer matrix ⁴. Denote the transfer matrix for bringing a particle of type a and rapidity θ through N particles and ending up with a particle of type b by $T_{ab}(\theta)$. We are here suppressing the dependence on the rapidities $\theta_1, \dots, \theta_N$ of the N particles as well as the q^N labels c_i for the species of the N particles before the scattering process and the q^N labels d_i for the N particles after the process (q is the number of different particle species). The components of $T_{ab}(\theta)$ can be written in terms of the S-matrix elements as

$$(T_{ab}(\theta))_{c_i}^{d_i} \equiv \sum_{\text{all } k} S_{ac_1}^{d_1 k_1}(\theta - \theta_1) S_{k_1 c_2}^{d_2 k_2}(\theta - \theta_2) \dots S_{k_{N-1} c_N}^{d_N b}(\theta - \theta_N) \quad (5.8)$$

Pictorially, this is much easier to digest; it is



where each intersection is an S-matrix element. Using the fact that all of our S-matrices satisfy $S_{ab}^{cd}(0) = -\delta_{ac}\delta_{bd}$, scatterings at zero relative rapidity permute the colliding particles (with a sign), it is seen that the matrix $\text{tr}_o T(\theta) \equiv \sum_a T_{aa}(\theta)$ satisfies

$$-\text{tr}_o T(\theta = \theta_k) = T(\theta_k | \theta_{k+1}, \dots, \theta_{k-1}), \quad (5.9)$$

the transfer matrices appearing in (5.2). We can thus concentrate on diagonalizing $\text{tr}_o T(\theta)$ for general θ and then set θ to the different θ_k at the end. The Yang-Baxter relation ensures that the $\text{tr}_o T(\theta)$ commute for different θ and thus can be simultaneously diagonalized for all θ by a θ -independent set of eigenvectors; only the eigenvalues depend on θ .

We do not need to find the eigenvalues of $\text{tr}_o T(\theta)$ explicitly in order to do the thermodynamics; rather, we derive constraint equations so that we can define eigenvalue densities just like the rapidity densities. This problem is equivalent to diagonalizing a transfer matrix in an integrable lattice model, where the S-matrix elements play the role of the lattice Boltzmann weights. There are a variety of techniques to do this [38]. One

⁴ We are grateful to N. Reshetikhin for this presentation.

which is particularly elegant and which is applicable to our models is the Algebraic Bethe Ansatz [39]. We will discuss this in appendix A. In the next section we apply these constraint equations to a variety of models. The result is that we will be able to obtain a TBA system of equations of the form (5.6)(5.7) containing some extra fictitious, zero-mass, particle species to account for the eigenvalue contribution to (5.2).

5.3. Casimir Energy in the UV Limit

The free energy (5.7) can be found explicitly in the $m_a R \rightarrow 0$ limit [40,4]. For zero chemical potential, it is given by

$$E(m_a R \rightarrow 0) \sim -\frac{1}{\pi R} \sum_a [\mathcal{L}(\frac{x_a}{1+x_a}) - \mathcal{L}(\frac{y_a}{1+y_a})], \quad (5.10)$$

where $\mathcal{L}(x)$ is Rogers dilogarithm function[41]

$$\mathcal{L}(x) = -\frac{1}{2} \int_0^x dy \left[\frac{\ln y}{(1-y)} + \frac{\ln(1-y)}{y} \right],$$

$x_a = \exp(-\epsilon(0))$, and $y_a = \exp(-\epsilon(\infty))$. The ground-state energy is found by setting all $\lambda_a = 1$. It follows from (5.6) that the constants x_a are the solutions to the equations

$$x_a = \prod_b (1 + \lambda_b x_b)^{N_{ab}}, \quad (5.11)$$

where $N_{ab} = \frac{1}{2\pi} \int d\theta \phi_{ab}(\theta)$. (We restore the chemical potentials because we will use them in sect. 6.3) The constants y_a in (5.10) are nonzero only for those species a' with $m_{a'} = 0$, where they are the solutions to

$$y_{a'} = \prod_{b'} (1 + \lambda_b y_{b'})^{N_{a'b'}}, \quad (5.12)$$

where b' also runs only over massless species. The UV limit Casimir energy (5.10) is to be compared with that of the conformal theory

$$E(R) = \frac{2\pi}{R} (h + \bar{h} - \frac{c}{12}), \quad (5.13)$$

where $(h, \bar{h})$ are the conformal dimensions of the operator which create this state. For unitary theories, the ground state has $h = \bar{h} = 0$ in (5.13). We can check a proposed scattering theory by comparing (5.10) with (5.13). Higher order in $m_a R$ corrections to (5.10) can be numerically computed from (5.6) and compared to perturbation theory about the UV limit conformal theory.

5.4. Casimir Energy in the IR limit

The Casimir energy in the IR limit can be expanded in a series in $e^{-m_a R}$. From (5.6) it is easy to show that for $m_a R$ large,

$$\epsilon_a(\theta) \approx m_a R \cosh \theta - \sum_{b'} N_{ab'} \ln(1 + \lambda_{b'} y_{b'}) \quad (5.14)$$

Plugging this into (5.7), one finds

$$E(mR \rightarrow \infty) \rightarrow - \sum_a q_a \int d\theta \frac{m_a}{2\pi} \cosh \theta e^{-m_a R \cosh \theta}, \quad (5.15)$$


where

$$q_a = \prod_{b'} (1 + \lambda_{b'} y_{b'})^{N_{ab'}}.$$

This provides a useful check on the results: q_a must be the number of particles of mass m_a because in the $R \rightarrow \infty$ limit we have a very dilute gas of particles, and the first term in this expansion of the free energy is given by one-particle contributions only. In the case where there are solitons, q_a is not required to be an integer: it is given by $B^{1/N}$, where B is the number of N -particle configurations.

6. TBA for Minimal Models Perturbed in the Least Relevant Operator

In this section we obtain and analyze the TBA system of equations for the $N=2$ scattering theories discussed in sect. 4. We will also consider the analogous $N=1$ case and the case of all perturbed $SU(2)_l \otimes SU(2)_k / SU(2)_{k+l}$ coset models, since this involves little additional work. From the form of the S-matrix (4.4) we see that the $N=0$ case enters as an ingredient for our $N=2$ case (and also for the $N=1$ case [19,20]). The RSOS S-matrix conjectured in [10] to describe the $N=0$ minimal models $SU(2)_1 \otimes SU(2)_k / SU(2)_{k+1}$ perturbed in the least relevant operator is believed to lead to a TBA system (5.6) which can be described by the figure [13]



Each node in the figure represents a particle species for the equations (5.6); the $k-1$ open nodes have $m_a = 0$ while the node labeled $\otimes$ is massive. The elements ϕ_{ab} in (5.6) are zero

unless the nodes associated with species a and b are connected by a line, in which case $\phi_{ab}(\theta) = (\cosh(\theta))^{-1}$. In other words, $\phi_{ab}(\theta) = l_{ab}\phi(\theta)$ where l_{ab} is the incidence matrix for the above diagram. This TBA system of coupled integral equations was analyzed in [13]; the system was proven to describe the RSOS S-matrix only for $k = 1$ and $k = 2$, but there is much evidence that the correspondence holds for the whole series. We shall assume that this is true in this section.

It is straightforward to take the UV and IR limits in the manner of sects. 5.3 and 5.4. With this ϕ_{ab} , we have $N_{ab} = \frac{1}{2}l_{ab}$. The solution to the equations (5.11) is [37,13]

$$x_a = \left(\frac{\sin \frac{\pi(a+1)}{k+3}}{\sin \frac{\pi}{k+3}} \right)^2 - 1. \quad (6.1)$$

To solve the equations (5.12), we notice that this case corresponds to the above Dynkin diagram with the k -th node removed. Thus the solution is

$$y_a = \left(\frac{\sin \frac{\pi(a+1)}{k+2}}{\sin \frac{\pi}{k+2}} \right)^2 - 1. \quad (6.2)$$

Plugging these into (5.10), and using the amazing dilogarithm identity [40,41]

$$\sum_{a=1}^k \mathcal{L} \left(\frac{\sin^2 \frac{\pi}{k+3}}{\sin^2 \frac{\pi(a+1)}{k+3}} \right) = \frac{2k}{k+3} \frac{\pi^2}{6} \quad (6.3)$$

along with $\mathcal{L}(1-x) = \mathcal{L}(1) - \mathcal{L}(x)$ and $\mathcal{L}(1) = 1$, yields

$$E(mR \rightarrow 0) \rightarrow - \left(\frac{\pi}{6R} \right) \left(1 - \frac{6}{(k+2)(k+3)} \right). \quad (6.4)$$

This, of course, is the correct UV limit. The first corrections to (6.4) also agree with the predictions from perturbed conformal field theory [13,6]. The infrared limit gives a nice check: the number of N -particle configurations for large N for the k -well soliton structure is given by q^N , where $q = 2 \cos \frac{\pi}{k+2}$ is the largest eigenvalue of the incidence matrix l_{ab} . Using the y_a in (5.15) gives precisely this result.

6.1. $N=2$ $X^3 - \beta X$

In this section, we apply the discussion of the appendix to obtain the TBA equations for our basic $N=2$ S-matrix $S_{k=1}^{N=2}$ (4.2). The eigenvalues of the N -particle transfer matrix are obtained from (A.10) to be

$$\lambda(\theta; \{y\}) = \mathcal{A} \prod_{j=1}^N Z(\theta - \theta_j) \prod_{r=1}^n \cosh\left(\frac{\theta - y_r}{2}\right) \prod_{r=n+1}^N i \sinh\left(\frac{\theta - y_r}{2}\right), \quad (6.5)$$

where $Z(\theta)$ is given by (4.3) and where we define

$$\mathcal{A} \equiv \prod_{r=1}^N (i \sinh(\frac{\theta - y_r}{2}))^{-1} (\prod_{j=1}^N i \sinh(\frac{\theta - \theta_j}{2}) + (-1)^n \prod_{j=1}^N \cosh(\frac{\theta - \theta_j}{2})). \quad (6.6)$$

The y_r in (6.5) are the N distinct solutions to (A.9), which here becomes

$$\prod_{j=1}^N \frac{i \sinh(\frac{y_r - \theta_j}{2})}{\cosh(\frac{y_r - \theta_j}{2})} = (-1)^{n+1}, \quad (6.7)$$

for any choice of $n = 1, \dots, N$. Note that, given (6.7), $\mathcal{A}$ has no poles in θ and is bounded at infinity and, thus, does not depend on θ .

Unitarity of the S-matrix means that the eigenvalues (6.5) have magnitude one, which is required for (5.2). The corresponding solutions to condition (6.7) are of the form $y = z_0 + i\pi/2$ and $y = z_{\bar{0}} - i\pi/2$, with z_0 and $z_{\bar{0}}$ real. Defining the distributions of such solutions as $P_0(z_0)$ and $P_{\bar{0}}(z_{\bar{0}})$ respectively, the log of (6.7) gives

$$2\pi P_i(z_i) = \int d\theta \frac{\rho(\theta)}{\cosh(z_i - \theta)}, \quad (6.8)$$

where $i = 0, \bar{0}$.

Condition (5.2) yields the relation

$$mL \cosh \theta + Im \frac{d}{d\theta} \ln \lambda(\theta) = 2\pi P(\theta) \quad (6.9)$$

for the distribution of allowed rapidities. Thus the level density for the real particles depends on the eigenvalue distributions, just as (5.4) depends on the rapidity distribution for all the particles. Each of the 2^N eigenvalues $\lambda(\theta, \{y\})$ is determined by specifying whether a given y_r contributes a $\cosh(\frac{1}{2}(\theta - y_r))$ or an $i \sinh(\frac{1}{2}(\theta - y_r))$ to the product in (6.5). (Remember that a given y_r appears exactly once in this product.) We thus define the density $P_i^-(z_i)$ of z_i which contribute a cosh to the product (n contributions in total), and $P_i^+(z_i)$ for the remaining contributions. We thus have the relation $P_i(z_i) = P_i^-(z_i) + P_i^+(z_i)$. With these definitions, (6.5) becomes

$$\begin{aligned} Im \frac{d}{d\theta} \ln \lambda(\theta) &= \int d\theta' \rho(\theta') Im \frac{d}{d\theta} \ln Z(\theta - \theta') \\ &+ \frac{1}{2} \sum_{i=1}^2 \int dz_i \frac{P_i^+(z_i) - P_i^-(z_i)}{\cosh(z_i - \theta)}, \end{aligned} \quad (6.10)$$

Now from (4.3), (6.8), and the integrals

$$\frac{1}{4} \int dt \frac{\cos \theta t}{\cosh^2(\frac{\pi t}{2})} = \int \frac{dy}{2\pi} \frac{1}{\cosh(y - \theta) \cosh y} = \frac{\theta}{\pi \sinh \theta},$$

we obtain

$$\int d\theta' \rho(\theta') \operatorname{Im} \frac{d}{d\theta} \ln Z(\theta - \theta') = \frac{1}{2} \sum_{i=1}^2 \int dz_i \frac{P_i(z_i)}{\cosh(z_i - \theta)}.$$


Relations (6.9) and (6.10) then become

$$mL \cosh \theta + \sum_{i=1}^2 \int dz_i \frac{P_i^+(z_i)}{\cosh(z_i - \theta)} = 2\pi P(\theta). \quad (6.11)$$

The relations (6.8) and (6.11) are constraints on the particle densities just like (5.4). Thus we can proceed identically to the diagonal case. We minimize the free energy with respect to ρ and P_i^+ , subject to the constraints (6.8) and (6.11). Defining

$$\frac{\rho(\theta)}{P(\theta)} = \frac{e^{-\epsilon(\theta)}}{1 + e^{-\epsilon(\theta)}} \quad \frac{P_i^+(z_i)}{P_i(z_i)} = \frac{e^{-\epsilon_i(z_i)}}{1 + e^{-\epsilon_i(z_i)}} \quad (6.12)$$

for $i = 0, \bar{0}$, it is seen that we obtain a TBA system of the form (5.6) and (5.7) consisting of three species, one massive and two massless, with $\phi_{ab}(\theta) = (\cosh \theta)^{-1} l_{ab}$, where l_{ab} is the incidence matrix for



The open nodes in the figure correspond to the massless pseudoparticles (the P_i^+ densities) and the solid node corresponds to the soliton (ρ density) with mass $m = |\Delta W|$.


To take the ultraviolet limit of this TBA system, we use our analysis of the non-supersymmetric case. Since l_{ab} is the incidence matrix for A_3 , we can read off the x_a from (6.1): $x_0 = x_{\bar{0}} = 2$ and $x_1 = 3$. The y_a can be found by removing the center node, leaving two unconnected nodes and $y_0 = y_{\bar{0}} = 1$. Using the dilogarithm identity (6.3) yields

$$E(mR \rightarrow 0) \rightarrow -\left(\frac{\pi}{6R}\right). \quad (6.13)$$

Thus our TBA system has the correct central charge $c = 1$ in the UV limit. The infrared limit also gives the correct result: we have 2^N different N -particle configurations.

6.2. $N=2$ minimal model with $\Phi_{(1,3)}$ perturbation

To find the TBA system for the entire $N=2$ series, we must diagonalize the transfer matrix (5.8) associated with the S-matrix (4.4). Since the S-matrix is a tensor product of an $N=0$ S-matrix with a supersymmetric one, we can diagonalize the two parts separately.⁵ Each diagonalization has already been done: the $N=0$ minimal-model part was discussed at the beginning of this section, while the supersymmetric part was done in the last subsection. Each piece contributes their associated pseudoparticles (the open nodes in the diagrams); the tensor-product structure means that the pseudoparticles from each piece do not couple to ones from the other. However, both types of pseudoparticle couple to the same massive particle, because both contribute to the single equation (6.9). Therefore, the TBA system has a single massive particle. It follows that the TBA equations are described by the $N=0$ figure for $S_k^{N=0}$ joined at the massive node to the basic $N=2$ part:



We do not know of any significance to the fact that this is a D_{k+2} Dynkin diagram.

We can now compute the UV limit of the Casimir energy. The x_a are given by [37]

$$\begin{aligned} x_0 &= x_{\bar{0}} = k + 1 \\ x_a &= (a + 1)^2 - 1 \end{aligned} \tag{6.14}$$

To find the y_a , we remove the massive node, leaving us with the A_{k-1} Dynkin diagram plus two individual nodes. Thus the y_a for $a = 1 \dots k$ are given by (6.2), while $y_0 = y_{\bar{0}} = 1$. There exists another amazing dilogarithm identity [40] (used in the D_n section of [37]) which gives us

$$E(mR \rightarrow 0) \rightarrow - \left(\frac{\pi}{6R} \right) \frac{3k}{k+2}, \tag{6.15}$$

in agreement with the central charge $\frac{3k}{k+2}$ for the $N=2$ minimal series. The IR limit is correct as well. It can also be shown from the TBA analysis that the bulk contribution to $E(R)$ (terms proportional to e.g. R or $R \ln R$ in the small R expansion) vanishes. This is to be expected since our theories are supersymmetric. We will now calculate $\text{tr}(-1)^F$ as another check on the supersymmetric nature of our theories.

⁵ We emphasize that the model is not a product of two models, since that would yield a direct product of S-matrices. In our situation, a particle can be thought of as a bound-state of a particle in the k -th minimal model with one in the $N=2$ $X^3 - X$ model.

6.3. $\text{tr}(-1)^F$

To calculate $\text{tr}(-1)^F$, we use the fact that with chemical potentials μ_a , the free energy is equal to $-T \ln \text{tr}(e^{\mu_a N_a/T})$. Thus in a theory with a diagonal S-matrix, one can calculate $\text{tr}(-1)^F$ by setting the chemical potentials $\mu_a = i\pi e_a T$, where e_a is the fermion number of a particle of species a . Finding the appropriate chemical potentials in a non-diagonal scattering theory is more subtle, because the diagonalization of the transfer matrix means that we are not working in a basis of particle eigenstates. However, the thermodynamic expectation value of a symmetry operator is well-defined on each state in this new basis because these operators commute with the Hamiltonian and hence the transfer matrix. Therefore, we need to find the eigenvalue of the symmetry operator associated with each eigenvalue of the transfer matrix. With some methods of determining transfer matrix eigenvalues, it is not obvious how to do this. For example, in the inversion-relation method used in [13], one finds the constraint equations for the eigenvalues without constructing the eigenstate. However, using the Algebraic Bethe Ansatz, we have an explicit expression (A.5) for the eigenstates, making this identification simple.

In our series of models, the particles (u, d) have fermion number $(\frac{1}{2}, -\frac{1}{2})$. Our eigenvector ψ (A.5) thus has fermion number $n - (N/2)$. This is the fermion number associated with each eigenvalue (6.5). By definition, we have

$$\begin{aligned} \int P_0^+ + \int P_{\bar{0}}^+ &= N - n \\ \int \rho &= N. \end{aligned} \tag{6.16}$$

To calculate $\text{tr}(-1)^F$, we set $\mu_0 R = \mu_{\bar{0}} R = -i\pi$, and $\mu_k R = \frac{i\pi}{2}$. The remaining chemical potentials for the pseudoparticles $1, \dots, (k-1)$ remain zero; these particles arise from diagonalizing the part of the S-matrix independent of fermion number.

To finish the calculation, we need to calculate the x_a and y_a from (5.11) and (5.12). The result is that with these chemical potentials, $x_a = y_a$. Specifically, $x_k = y_k = 0$, $x_0 = y_0 = x_{\bar{0}} = y_{\bar{0}} = 1$, and the remaining ones are as in (6.2). The fact that $x_a = y_a$ means that $E_\lambda(R) = 0$ for all R —not only in the UV limit. Thus

$$\text{tr}(-1)^F = e^{-E(R)L} = 0 + \mathcal{O}(1). \tag{6.17}$$

Since we are working in the large L limit, we have proven only that the leading term, proportional to L , vanishes. This, of course, is required in a supersymmetric theory—(6.17) gives the number of bosonic ground states minus the number of fermionic ones,

because the others pair up and cancel. (For the k theory, $\text{tr}(-1)^F = k + 1$.) Thus this calculation gives no new information, but does provide a check on our TBA system, as well as giving as simple example of the application of the excited-state TBA analysis [7] to models with pseudoparticles.


6.4. $N=1$ minimal models with $\Phi_{(1,3)}$ superfield perturbation

The S-matrix for the scattering theory coming from the $N=1$ minimal model $SU(2)_2 \otimes SU(2)_k / SU(2)_{k+2}$ perturbed in the least relevant, supersymmetry preserving operator (the $\Phi_{(1,3)}$ superfield) was conjectured in [19,20] to be

$$S_k^{N=1}(\theta) = S_k^{N=0}(\theta) \otimes S_{k=1}^{N=1}(\theta), \quad (6.18)$$

where $S_{k=1}^{N=1} = S_{k=2}^{N=0}$, the S-matrix for the perturbed tricritical Ising model. This was conjectured because it is the simplest S-matrix consistent with the appropriate quantum-group symmetry and the supersymmetry [20]. The first part of this tensor product arises because the Landau-Ginzburg picture [15] predicts solitons interpolating between the minima of a Chebyshev superpotential, just as in the $N=2$ case. The second part gives the model its supersymmetric structure, which acts nonlocally on solitons [10]. The origin of this soliton structure is unclear. Presumably there is a bosonization of this series analogous to the sine-Gordon description of the $N=2$ $X^3 - X$ model, and the solitons interpolate between vacua of this field.

To find the TBA system, we must diagonalize the transfer matrix (5.8) associated with the S-matrix (6.18). As in the $N=2$ series, the S-matrix is a tensor product, and we diagonalize the two parts separately. Each piece is an $N=0$ minimal model, so this diagonalization has already been done [13], and is discussed at the beginning of this section. As in the $N=2$ series, the resulting two sets of pseudoparticles are coupled through the massive node:



It is simple to take the UV limit of this system using the analysis of the $N=0$ minimal models. Since we now have $k + 1$ nodes instead of k , the x_a are given by (6.1) with the substitution $k \rightarrow (k + 1)$. When the massive node is removed, we obtain a diagram of

$k - 1$ nodes along with a single disconnected node. Thus the y_a for $a = 1 \dots k$ are given by (6.2), while $y_{k+1} = 1$. Using the dilogarithm identity (6.3) again, we find that

$$E(mR \rightarrow 0) \rightarrow - \left(\frac{\pi}{6R} \right) \left(\frac{3}{2} - \frac{12}{(k+2)(k+4)} \right), \quad (6.19)$$

in agreement with the conformal results.

We note that the $k = 2$ model in this series corresponds to the first model in the $N=2$ series. However, the S-matrix (6.18) for $k = 2$ is not identical to the S-matrix (4.1). In fact, the particle content is different: the particle content here is not the soliton doublet discussed above but instead is that of the tensor product of two triple-well soliton potentials. This is equivalent to making the vacua the nodes and the solitons the links of the Dynkin diagram for $\hat{A}_3$ or $\hat{D}_4$, depending on how one does the tensor product. However, one can recover the S-matrix elements (4.1) from (6.18) by taking an orbifold, where one mods out by a symmetry. The orbifold procedure is a generalization of the Kramers-Wannier high- to low-temperature duality. This redefines the states as linear combinations of the previous states, hence changing the S-matrix. The new S-matrix can be found in the manner of lattice-model orbifolds [43], and one recovers the S-matrix (4.1). This was done in [10] to relate an $N=0$ D -series S-matrix (the tricritical three-state Potts model) to one in the A -series. In fact, this can be done for every other model in the D -series, as in the lattice models of [43]. The orbifolding should not change the ground-state energy, but will affect excited-state energies [44].

6.5. $SU(2)_l \otimes SU(2)_k / SU(2)_{k+l}$

The general $SU(2)$ coset case is the obvious generalization. The S-matrix here is [20]

$$S_{k,l}^{N=0}(\theta) = S_k^{N=0}(\theta) \otimes S_l^{N=0}(\theta), \quad (6.20)$$

It follows that the TBA system is described by the diagram

$$\begin{array}{ccccccccc} 1 & 2 & & k-1 & k & & k+1 & & k+l-1 \\ \bigcirc & \text{---} & \bigcirc & \text{---} & \bigcirc & \text{---} & \bigotimes & \text{---} & \bigcirc & \text{---} & \bigcirc \end{array}$$

The x_a are given by (6.1) with the substitution $k \rightarrow k + l - 1$. The y_a for $a = 1, \dots, k$ are given by (6.2), while for $a = k + 1, \dots, k + l - 1$ they are given by (6.2) with $a \rightarrow a - k$ and $k \rightarrow l$. The result for the energy is in agreement with the central charge of the conformal theory $c = 3(\frac{k}{k+2} + \frac{l}{l+2} - \frac{k+l}{k+l+2})$.

7. Zamolodchikov's c-function and more

The ground state energy $E(R)$ described in the previous section in terms of the solutions to our set of coupled integral equations can be thought of as a Zamolodchikov 'c-function' [45] for our massive theory: $E(R) = -\pi c(mR)/6R$. It is not obvious from the TBA equations that this 'c-function' monotonically decreases in the infra-red, though this is observed to be the case in the numerical and analytic analysis of all physical TBA systems checked⁶.

In [29] another 'c-function' was obtained for the Chebyshev theories using the $N=2$ structure. It is interesting to compare these results. The c-function obtained in [29] is quite simple:

$$c_k(mR = z) = \frac{3}{2}z \frac{d}{dz} u(z, r = 2k/(k+2)) \quad (7.1)$$

where $u(z, r)$ is a solution of the differential equation $u_{zz} + z^{-1}u_z = \sinh u$ (which can be transformed to Painleve III). The boundary conditions of u are specified by (for $r < 2$)

$$u(z, r) \sim r \log z + \text{const.} + \dots \quad \text{as } z \rightarrow 0. \quad (7.2)$$

The values for r in the solutions (7.1) are obtained from regularity conditions. The only k dependence is in the boundary conditions of u . It is easily seen from (7.2) that (7.1) yields the correct central charges in the UV limit $z \rightarrow 0$. It would be interesting to directly calculate the next UV corrections to the 'c-function' from perturbation theory around the conformal theory and compare with that predicted by the TBA approach and that predicted by (7.1). Comparing the IR limits ($z \rightarrow \infty$), it can be seen that the two approaches give different results: the c-function (7.1) has contributions corresponding to only odd numbers of particles, whereas the TBA approach includes contributions from any number of particles. It remains to be seen if one can relate these approaches by some projection in the manner of [7]. If this were possible it would be quite interesting: we could work with an differential equation (and a well-studied one at that) rather than the TBA system of coupled integral equations (5.6).

⁶ T.R. Klassen and E. Melzer, private communication

8. Conclusions

We have found the exact S-matrix for the $N=2$ discrete series perturbed by the least-relevant supersymmetry-preserving operator. The nice thing about the $N=2$ non-renormalization theorems is that they leave little guesswork; here one does not have to conjecture the particle content. Since our results fit in so nicely with the $N=0$ discrete series, we believe that this lends more support to the conjectured $N=0$ soliton picture, and that much intuition can be gained by using the $N=2$ models to study the $N=0$ models.

Another interesting aspect is the appearance of solitons with fractional fermion number: $\pm\frac{1}{2}$ here, and any rational number in our next paper. There are only a few other integrable models with such structure, and those have only fractional charge $\pm\frac{1}{2}$ [46]. Therefore, this analysis may be valuable for studying experimentally-realizable systems of fractional fermion number. It is probably not possible to measure the S-matrix or the Casimir energy directly, but one knows many other things about these $N=2$ systems which may be observable. For example, all the critical exponents of these systems are known exactly.

Acknowledgements

We are very grateful to N. Reshetikhin for many helpful conversations on the TBA, and for providing the presentation of sect. 5.2. We would also like to thank C. Vafa for valuable discussions, and T. Klassen and E. Melzer for their comments on the manuscript. K.I. was supported by NSF grant PHY-87-14654 and an NSF graduate fellowship, and P.F. was supported by DOE grant DEAC02-89ER-40509.

Appendix A. Algebraic Bethe Ansatz for the 6-vertex Model

Consider a scattering theory consisting of two species u and d with the two particle S-matrix:

$$\begin{array}{cc} du & ud \\ \begin{array}{c} ud \\ du \end{array} \left(\begin{array}{cc} b & \tilde{c} \\ c & \tilde{b} \end{array} \right) & \begin{array}{cc} uu & dd \\ uu & dd \end{array} \left(\begin{array}{cc} a & 0 \\ 0 & \tilde{a} \end{array} \right). \end{array} \quad (\text{A.1})$$

Our S-matrix elements are taken to satisfy the condition

$$a(\theta)\tilde{a}(\theta) + b(\theta)\tilde{b}(\theta) - c(\theta)\tilde{c}(\theta) = 0. \quad (\text{A.2})$$

Drawing a scattering vertex with right or up pointing arrows to denote species u and left or down arrows to denote species d , the scattering theory (A.1) corresponds to the 6-vertex model. Condition (A.2) is known as the free fermion condition in the six-vertex model[38]. We will discuss the algebraic Bethe ansatz technique for finding the eigenvalues of the transfer matrix for theory (A.1). Note that our basic $N=2$ scattering theory is of this type; we are here being more general with future applications in mind.

The matrix $T_{ab}(\theta)$ discussed in sect. 4.2 will be denoted by

$$T_{ab}(\theta) = \begin{pmatrix} A(\theta) & B(\theta) \\ C(\theta) & D(\theta) \end{pmatrix}; \quad (\text{A.3})$$

$C(\theta)$, for example, is the transfer matrix for bringing a particle of type d and rapidity θ through N particles with rapidities θ_i and ending up with a particle of type u coming out. As discussed in sect. 4.2, we will be interested in finding the eigenvalues of the $2^N \times 2^N$ matrix $tr_o T = A(\theta) + D(\theta)$.

Define $\Omega \equiv |\prod_{i=1}^N d(\theta_i)\rangle$. Notice from (A.1) that for any rapidities we have $C(\theta)\Omega = 0$. The state Ω is also an eigenstate of the matrices A and D :

$$A(\theta)\Omega = \prod_{i=1}^N b(\theta - \theta_i)\Omega, \quad D(\theta)\Omega = \prod_{i=1}^N \tilde{a}(\theta - \theta_i)\Omega. \quad (\text{A.4})$$

The algebraic Bethe ansatz is that all eigenvectors of the matrix $A + D$ can be written as

$$\psi = \prod_{r=1}^n B(y_r)\Omega, \quad (\text{A.5})$$

for some choices of the y_r and n (it follows from (A.1) that the vector (A.5) has n u particles and $N - n$ d particles). As a result of the Yang-Baxter relation, $T_{ab}(\theta)$ satisfies the relation

$$S_{ac}^{a'c'}(\theta - y_r)T_{a'b}(\theta)T_{c'd}(y_r) = T_{cd'}(y_r)T_{ab'}(\theta)S_{b'd'}^{db}(\theta - y_r) \quad (\text{A.6})$$

This yields the following useful commutation relations for the transfer matrices:

$$\begin{aligned} A(\theta)B(y_r) &= \frac{a(\theta - y_r)}{b(\theta - y_r)}B(y_r)A(\theta) - \frac{c(\theta - y_r)}{b(\theta - y_r)}B(\theta)A(y_r) \\ D(\theta)B(y_r) &= -\frac{a(\theta - y_r)}{b(\theta - y_r)}B(y_r)A(\theta) + \frac{c(\theta - y_r)}{b(\theta - y_r)}B(\theta)D(y_r), \end{aligned} \quad (\text{A.7})$$

provided $b(\theta - y_r) \neq 0$. We have used the condition (A.2) to simplify the equations. It follows from these relations and (A.4) that $A(\theta) + D(\theta)$ act on the states (A.5) as

$$[A(\theta) + D(\theta)]\psi = \prod_{r=1}^n \frac{a(\theta - y_r)}{b(\theta - y_r)} \left(\prod_{i=1}^N b(\theta - \theta_i) + (-1)^n \prod_{i=1}^N \tilde{a}(\theta - \theta_i) \right) \psi + \text{bad terms}, \quad (\text{A.8})$$

where the ‘bad terms’ are vectors related to (A.5) by some $B(y_r) \rightarrow B(\theta_p)$. All such bad terms are proportional to $[A(y_r) + (-1)^n D(y_r)]\Omega$ and thus vanish if

$$\prod_{i=1}^N \frac{b(y_r - \theta_i)}{\tilde{a}(y_r - \theta_i)} = (-1)^{n+1}, \quad (\text{A.9})$$

this determines the y_r . Actually, since our S-matrices all have $b(0) = 0$, the argument (using (A.7)) that all bad terms are proportional to (A.9) breaks down if some of the y_r are the same. We thus require the solutions of (A.9) to have no repeating values of y_r . Our desired eigenvalues of $\text{tr}_o T(\theta)$ are thus given by

$$\lambda(\theta; y) = \prod_{r=1}^n \frac{a(\theta - y_r)}{b(\theta - y_r)} \left(\prod_{i=1}^N b(\theta - \theta_i) + (-1)^n \prod_{i=1}^N \tilde{a}(\theta - \theta_i) \right), \quad (\text{A.10})$$

for any choice of distinct solutions y_r and n of (A.9).

References

- [1] A.B. Zamolodchikov and Al.B. Zamolodchikov, *Ann. Phys.* 120 (1980) 253.
- [2] A.B. Zamolodchikov, *Adv. Stud. Pure Math.* 19 (1989) 1.
- [3] C.N. Yang and C.P. Yang, *J. Math. Phys.* 10 (1969) 1115.
- [4] Al.B. Zamolodchikov, *Nucl. Phys.* B342 (1990) 695.
- [5] M.J. Martins, *Phys. Rev. Lett.* 67 (1991) 419.
- [6] T. Klassen and E. Melzer, “Spectral Flow between Conformal Field Theories in 1+1 Dimensions”, Chicago preprint EFI 91-17, Miami preprint UMTG-162.
- [7] P. Fendley, “Excited-state thermodynamics”, Boston preprint BUHEP-91-16.
- [8] A.B. Zamolodchikov, *JETP Lett.* 46 (1987) 160.
- [9] Al.B. Zamolodchikov, *Nucl. Phys.* B358 (1991) 524.
- [10] A.B. Zamolodchikov, Landau Institute preprint, September 1989.
- [11] A. LeClair, *Phys. Lett.* 230B (1989) 103.
- [12] H. Itoyama and P. Moxhay, *Phys. Rev. Lett.* 65 (1990) 2102.
- [13] Al.B. Zamolodchikov, *Nucl. Phys.* B358 (1991) 497.
- [14] P. Fendley, W. Lerche, S.D. Mathur and N.P. Warner, *Nucl. Phys.* B348 (1991) 66;
P. Mathieu and M. Walton, *Phys. Lett.* 254B (1991) 106.
- [15] A.B. Zamolodchikov, *Sov. J. Nucl. Phys.* 44 (1986) 529;
D. Kastor, E. Martinec and S. Shenker, *Nucl. Phys.* B316 (1989) 590;
E. Martinec, *Phys. Lett.* 217B (1989) 431;
C. Vafa and N.P. Warner, *Phys. Lett.* 218B (1989) 51.
- [16] D. Bernard and A. LeClair, *Phys. Lett.* 247B (1990) 309.
- [17] P. Fendley, S.D. Mathur, C. Vafa and N.P. Warner, *Phys. Lett.* B243 (1990) 257.
- [18] G.E. Andrews, R.J. Baxter and P.J. Forrester, *J.Stat.Phys.* 35 (1984) 193.
- [19] K. Schoutens, *Nucl. Phys.* B344 (1990) 665.
- [20] C. Ahn, D. Bernard and A. LeClair, *Nucl. Phys.* B336 (1990) 409
- [21] W. Lerche and N.P. Warner, *Nucl. Phys.* B358 (1991) 571.
- [22] R. Jackiw and C. Rebbi, *Phys. Rev.* D13 (1976) 3398.
- [23] P. Fendley and K. Intriligator, “Son of Scattering and Thermodynamics of Fractionally-Charged Supersymmetric Solitons”.
- [24] E. Witten and D. Olive, *Phys. Lett.* 78B (1978) 97.
- [25] J. Goldstone and F. Wilczek, *Phys. Rev. Lett.* 47 (1981) 986
- [26] A.J. Niemi and G.W. Semenoff, *Phys. Rep.* 135 (1986) 99
- [27] W.P. Su, J.R. Schrieffer, A.J. Heeger, *Phys. Rev. Lett.* 42 (1979) 1698; *Phys. Rev.* B22 (1980) 2099.
W.P. Su and J.R. Schrieffer, *Phys. Rev. Lett.* 46 (1981) 738.
- [28] R. Dijkgraaf, E. Verlinde, and H. Verlinde, *Nucl. Phys.* B352 (1991) 59.

- [29] S. Cecotti and C. Vafa, “Topological Anti-Topological Fusion” Harvard and Trieste preprint HUTP-91/A033, SISSA-69/91/EP.
- [30] D. Gepner, *Fusion Rings and Geometry* preprint NSF-ITP-90-184.
- [31] W. Lerche and N. Warner, “Solitons in Integrable $N=2$ Supersymmetric Landau-Ginzburg Models”, to appear in the Proceedings of the Stony Brook conference “Strings and Symmetries”.
- [32] N. Reshetikhin and F. Smirnov, Commun. Math. Phys. 131 (1990) 157.
- [33] D. Bernard and A. LeClair, to appear in Commun. Math. Phys.
- [34] D. Friedan, Z. Qiu and S. Shenker, Phys. Lett. 151B (1985) 37;
Z. Qiu, Nucl. Phys. B270 (1986) 205.
- [35] N. Reshetikhin, J. Phys. A24 (1991) 3299.
- [36] V. Pasquier, Nucl. Phys. B285 (1987) 162.
- [37] T.R. Klassen and E. Melzer, Nucl. Phys. B338 (1990) 485; Nucl. Phys B350 (1990) 635.
- [38] R. Baxter, *Exactly Solved Models in Statistical Mechanics* Academic Press, 1982.
- [39] E.K. Sklyanin, L.A. Takhtadzhyan and L.D. Faddeev, Theo. Math. Phys. 40 (1980) 688. L. Faddeev, in Les Houches 1982 *Recent advances in field theory and statistical mechanics* (North-Holland 1984), edited by J.-B. Zuber and R. Stora.
- [40] A.N. Kirillov and N. Yu. Reshetikhin, J. Phys. A20 (1987) 1587.
- [41] L. Lewin, *Polylogarithms and Associated Functions* (North-Holland, 1981).
- [42] H.W.J. Blöte, J.L. Cardy and M.P. Nightingale, Phys. Rev. Lett. 56 (1986) 742;
I. Affleck, Phys. Rev. Lett., 56 (1986) 746.
- [43] P. Fendley and P. Ginsparg, Nucl. Phys. B324 (1989) 549.
- [44] T.R. Klassen and E. Melzer, “RG Flows in the D series of Minimal CFTs”, Cornell preprint CLNS-91-1111.
- [45] A.B. Zamolodchikov, JETP Lett. 43 (1986) 730;
A. Ludwig and J.L. Cardy, Nucl. Phys. B285 (1987) 687.
- [46] R. Shankar and E. Witten, Nucl. Phys. B141 (1978) 349;
N. Andrei and C. Destri, Nucl. Phys. B231 (1984) 445.